

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea. **Causes serious eye irritation. Very toxic to aquatic life with long lasting effects.** Wear protective gloves / protective clothing/eye protection/face protection. If eye irritation persists: Get medical advice/attention. Collect spillage. Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty containers which can be disposed of as non-hazardous waste. Contains disodium maleate. May produce an allergic reaction. To avoid risks to human health and the environment, comply with the instructions for use.

WARNING



aldor[®]
FLEX

H

HERBICIDE

A herbicide for the pre-emergence and early post-emergence control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces (railway ballast only) and amenity vegetation (around).

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea.

MAPP 19033

10 g e

GB86263572C

ARTICLE 85374958 - P2

For 24 hour emergency information contact Bayer CropScience Limited
Telephone: 00800 1020 3333 or nearest National Poisons Information Centre

Bayer CropScience Limited
230 Cambridge Science Park, Milton Road
Cambridge, CB4 0WB
Telephone: 00800 1214 9451

www.environmentalscience.bayer.co.uk for SDS and larger label

© Registered trademark of Bayer CropScience Ltd.

Bayer



VALDOR® FLEX
 A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea
Causes serious eye irritation.
Very toxic to aquatic life with long lasting effects.
 Wear protective gloves / protective clothing/eye protection/face protection.
 If eye irritation persists: Get medical advice/attention.
 Collect spillage.
 Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty containers which can be disposed of as non-hazardous waste.
 Contains disodium maleate. May produce an allergic reaction.
 To avoid risks to human health and the environment, comply with the instructions for use.

Directions for use - FOR USE AS A PROFESSIONAL HERBICIDE
 Situation: For use on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, and hard surfaces (railway ballast only) and amenity vegetation (around).
 Maximum individual dose: 1 sachet (10 g) / 200 m²
 Maximum number of treatments: 1 per year.
 Aquatic buffer zone distance: Hand-held use with coarse nozzles – 2 metres.
 Vehicle-mounted use with three-star nozzles – 6 metres.

Other specific restrictions:
 This product must only be applied to natural or permeable surfaces such as gravel or railway ballast. Do not apply to any non-permeable man-made surfaces. Use on hard surfaces refers to railway ballast only.
 For hand-held use: To minimise spray drift, the product must be applied using a nozzle capable of producing a coarse quality spray. For horizontal boom sprayer use: Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Division's website. These operating conditions must be maintained until the operator is 30m from the top of the bank of any surface water bodies. Buffer zones greater than 5m are NOT eligible for buffer zone reduction under the LERAP scheme.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

HERBICIDE

A herbicide for the pre-emergence and early post-emergence control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces (railway ballast only) and amenity vegetation (around).

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea.

MAPP 19033

10 x 10 g e

Pre and early-post emergent

SAFETY PRECAUTIONS

Operator Protection
 Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
 WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the product.
 However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.
 WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.
 IF SWALLOWED, seek medical advice immediately and show this container or label.

Environmental Protection
 Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains, farmyards and roads).
 Extreme care must be taken to avoid spray drift onto non-target plants outside the target area.

Hand-held use: Since there is a risk to aquatic life from use, direct spray must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Vehicle-mounted boom sprayer: To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies as specified for the crop. HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY. Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Division's website. Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form. The LERAP record form must be kept available for three years.

Storage and Disposal
 KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
 KEEP OUT OF REACH OF CHILDREN.
 STORE UNUSED SACHETS in a safe place.
 DO NOT STORE half-used sachets.
 EMPTY CONTAINER COMPLETELY and dispose of safely.
 PROTECT FROM FROST.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

GB86267500C - ARTICLE 85374958 - P2

For 24 hour emergency information contact Bayer CropScience Limited
 Telephone: 00800 1020 3333 or nearest National Poisons Information Centre

Bayer CropScience Limited
 230 Cambridge Science Park
 Milton Road
 Cambridge, CB4 0WB
 Telephone: 00800 1214 9451
 www.environmentalscience.bayer.co.uk for SDS and larger label

© Registered trademark of Bayer CropScience Ltd.



Bayer

ARTICLE 85374958



valdor[®]
FLEX


HERBICIDE

A herbicide for the pre-emergence and early post-emergence control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces (railway ballast only) and amenity vegetation (around).

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea

MAPP 19033

12 x 10 x 10 g e



VALDOR[®] FLEX

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea.



WARNING



Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves / protective clothing/eye protection/face protection.

If eye irritation persists: Get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty containers which can be disposed of as non-hazardous waste.

Contains disodium maleate. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

For 24 hour emergency information contact Bayer CropScience Limited
Telephone: 00800 1020 3333 or nearest National Poisons Information Centre

GB86209306C - P2

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge, CB4 0WB
Telephone: 00800 1214 9451

www.environmentalscience.bayer.co.uk for SDS and larger label

© Registered trademark of Bayer CropScience Ltd.



2 500037 105559 2

Bayer



A herbicide for the pre-emergence and early post-emergence control of annual and perennial weeds on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces (railway ballast only) and amenity vegetation (around).

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea.

MAPP 19033

Valdor Flex is to be dispersed in water (1 sachet in 6 – 10 L) and should be applied using hand-held applicators or vehicle mounted sprayer. A drift shield is recommended for use around amenity vegetation, and in other areas where desired vegetation may be present.

Valdor Flex will not provide control of established deep rooted perennial weeds such as dandelion, thistle, dock and nettles.

RESTRICTIONS

Since there is a risk to aquatic life from use, direct spray from hand-held equipment must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Do not allow direct spray from horizontal boom sprayer to fall within 6 m of the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Applications should not be made to plants growing under stress conditions, such as drought or waterlogging, as reduced levels of control may result.

Do not spray in windy weather.

Extreme care must be taken to avoid drift onto non-target plants, this includes: all green plant parts such as leaf surfaces, young bark or suckers of valued plants. Failure to do so may result in permanent damage or plant death.

Where Valdor Flex has been applied to sites that are subsequently to be cleared or grubbed, a period of at least 6 months should elapse between treatment and the sowing and planting of subsequent crops. In addition, the soil should be deeply cultivated or dug afterwards to ensure thorough mixing in order to remove any risk of damaging subsequent crops or planting.

Where Valdor Flex or other products containing diflufenican are applied in successive years, levels of diflufenican will build up in the soil. Even where soils are thoroughly dug there may be a risk of damage to subsequent plantings.

Care should be taken when applying Valdor Flex as heavy rain following application may wash the herbicide onto sensitive areas such as newly sown grass and areas about to be planted.

Where the soil organic matter content is greater than 10%, or for example where leaves have been collected or where a mat of organic matter has built up, pre-emergence activity will be reduced.

For maximum persistence of activity the area treated should not be cultivated or raked following application.

For maximum pre-emergence and residual activity from Valdor Flex please ensure good coverage of the spray swath. Overdosing should be avoided.

DO NOT APPLY VALDOR FLEX OVER DRAINS OR IN DRAINAGE CHANNELS, GULLIES OR SIMILAR STRUCTURES FOR MOVING WATER.

WEEDS CONTROLLED

Strains of some annual weeds (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group (WRAG) and copies are available from the AHDB, CPA, your distributor, crop adviser or product manufacturer.

The presence of populations resistant to ALS-inhibiting herbicides may lead to unacceptable levels of control. To reduce the risk of developing resistance or where resistance to sulfonylurea herbicides is suspected, applications should be made to young, actively growing weeds.

Key aspects of the Valdor Flex resistance management strategy are:

- ALWAYS follow WRAG guidelines for preventing and managing herbicide resistant grass and broad-leaved weeds.
- IDEALLY apply Valdor Flex pre-emergence or to young actively growing weeds up to the 2 leaf stage.
- DO NOT use Valdor Flex as the sole means of grass weed or broad-leaved weed control in successive years.
- ALWAYS rotate use of grass and broad-leaved weed herbicides with non-ALS modes of action.
- ALWAYS monitor weed control effectiveness and investigate any odd patches of poor grass or broad-leaved weed control. If unexplained contact your agronomist or technical advisor, who may consider a resistance test appropriate.
- Only one application of Valdor Flex may be made per year.

Valdor Flex controls susceptible annual and perennial weeds if applied pre- or early post-emergence up to the 2-leaf stage of the weeds, as directed in **Rates of Use**. Effectiveness using three star drift reduction technology may be reduced.

On natural surfaces not intended to bear vegetation, permeable surfaces overlying soil and amenity vegetation (around)

Grass weeds:

- Annual meadow-grass (*Poa annua*)
- Cocksfoot (*Dactylis glomerata*)
- Perennial rye-grass from seed (*Lolium perenne*)

Broad-leaved weeds:

- Black nightshade (*Solanum nigrum*)
- Bristly oxtonge (*Helminthotheca echinoides*)
- Canadian fleabane (*Erigeron canadensis*)
- Common field speed-well (*Veronica persica*)
- Common groundsel (*Senecio vulgaris*)
- Common purslane (*Portulaca oleracea*)
- Common Stork's-bill (*Erodium cicutarium*)
- Cut-leaved crane's-bill (*Geranium dissectum*)
- Dandelion (*Taraxacum officinale*)
- Dove's-foot crane's-bill (*Geranium molle*)
- Fat hen (*Chenopodium album*)
- Field pansy (*Viola arvensis*)
- Greater plantain (*Plantago major*)
- Hairy bitter-cress (*Cardamine hirsuta*)
- Knotgrass (*Polygonum aviculare*)
- Lesser trefoil (*Trifolium dubium*)
- Mayweeds (*Matricaria* sp.)
- Mouse-ear hawkweed (*Pilosella officinarum*)
- Narrow-leaved ragwort (*Senecio inaequidens*)
- Perennial sow-thistle (*Sonchus arvensis*)
- Prickly sow-thistle (*Sonchus asper*)
- Ribwort Plantain (*Plantago lanceolata*)
- Rosebay willowherb (*Chamerion angustifolium*)
- Scarlet pimpernel (*Anagallis arvensis*)
- Shepherd's purse (*Capsella bursa-pastoris*)
- Smooth sow-thistle (*Sonchus oleraceus*)
- Sowthistles (*Sonchus* sp.)
- Spotted spurge (*Chamaesyce maculata*)
- Tussock hawkweed (*Hieracium lachenalii*)
- White clover (*Trifolium repens*)
- Willowherbs (*Epilobium* sp.)
- Yarrow (*Achillea millefolium*)

On hard surfaces (railway ballast only)

Grass weeds:

- Annual meadow-grass (*Poa annua*)

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the product.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

IF SWALLOWED, seek medical advice immediately and show this container or label.

Environmental Protection

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains, farmyards and roads). Extreme care must be taken to avoid spray drift onto non-target plants outside the target area.

Hand-held use: Since there is a risk to aquatic life from use, direct spray must not be allowed to fall within 2 m of the top of the bank of any static or flowing waterbody or the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Vehicle-mounted boom sprayer: To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies as specified for the crop. HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY. Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Division's website. Maintain three star operating conditions until 30 m from the top of the bank of any surface water bodies.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.

The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form. The LERAP record form must be kept available for three years.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

STORE UNUSED SACHETS in a safe place.

DO NOT STORE half-used sachets.

EMPTY CONTAINER COMPLETELY and dispose of safely.

PROTECT FROM FROST.

VALDOR® FLEX

A water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium, a sulfonylurea.



WARNING

Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves / protective clothing/eye protection/face protection.

If eye irritation persists: Get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty containers which can be disposed of as non-hazardous waste.

Contains disodium maleate. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

Directions for use

FOR USE AS A PROFESSIONAL HERBICIDE

Situation: For use on natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, hard surfaces (railway ballast only) and amenity vegetation (around).

Maximum individual dose: 1 sachet (10 g) / 200 m²

Maximum number of treatments: 1 per year.

Aquatic buffer zone distance: Hand-held use with coarse nozzles – 2 metres.

Vehicle-mounted use with three-star nozzles – 6 metres.

Other specific restrictions:

This product must only be applied to natural or permeable surfaces such as gravel or railway ballast.

Do not apply to any non-permeable man-made surfaces.

Use on hard surfaces refers to railway ballast only.

For hand-held use: To minimise spray drift, the product must be applied using a nozzle capable of producing a coarse quality spray.

For horizontal boom sprayer use: Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Division's website. These operating conditions must be maintained until the operator is 30m from the top of the bank of any surface water bodies. Buffer zones greater than 5m are NOT eligible for buffer zone reduction under the LERAP scheme.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

Valdor Flex is a water dispersible granule formulation containing 360 g/kg diflufenican and 10 g/kg iodosulfuron-methyl-sodium. Valdor Flex is a pre- and early post-emergence herbicide for control of a wide range of annual and perennial grasses and broad-leaved weeds for up to 4 months on non-crop areas (permeable surfaces overlying soil) such as: gravel paths and driveways, porous surfaces alongside roadways and fence lines, porous strips of land adjacent to buildings, industrial sites, timber yards, farm yards, oil and gas storage sites, power stations, electric sub-stations, beneath pylons, around street/park obstacles and furniture, porous areas near to trees and shrubs and other natural surfaces where vegetation is not desirable, and railway ballast. Valdor Flex may also be used around the base of trees, shrubs and other plants in areas of semi-natural or ornamental vegetation, including parks, roadsides and other amenity areas.

Valdor Flex is applied to give pre- and early post-emergence weed control up to the 2-leaf stage of the weeds. One application of Valdor Flex can be made per year. Valdor Flex can be applied during cold weather, however application to frozen ground should be avoided.

Broad-leaved weeds:

Bristly ox-tongue (*Helminthotheca echioides*)
Creeping thistle (*Cirsium arvense*)
Cut-leaved cranesbill (*Geranium dissectum*)
Dandelion (*Taraxacum officinale*)
Dove's-foot cranesbill (*Geranium molle*)
Field bindweed (*Convolvulus arvensis*)
Lesser trefoil (*Trifolium dubium*)
Mayweeds (*Matricaria* sp.)
Ribwort (*Plantago lanceolata*)
Rosebay willowherb (*Chamerion angustifolium*)
Shepherd's purse (*Capsella bursa-pastoris*)
Sow-thistles (*Sonchus* sp.)
Willowherbs (*Epilobium* sp.)

Well-developed or established weeds (greater than 2 true leaves) will not be controlled.

SUSCEPTIBILITY OF NON-TARGET SPECIES

Trials have been conducted to evaluate the susceptibility of ornamental plants which could be exposed to spray drift during application. The following deciduous trees, shrubs and conifer species are resistant to the product when applied as recommended. Transient effects such as discoloration or chlorosis may occur if spray drift comes in to direct contact with the foliage, but this should have no long-lasting adverse effect on the plants.

Alder (*Alnus glutinosa*)
American alder (*Alnus incana*)
American red oak (*Quercus rubra* L.)
Ash-leaved maple (*Acer negundo*)
Bull bay (*Magnolia grandiflora*)
Canoe birch (*Betula papyrifera*)
Common rowan (*Sorbus aucuparia*)
Crab apple (*Malus sylvestris*)
Elms (*Ulmus* L. spec.)
English oak (*Quercus robur*)
European ash (*Fraxinus excelsior*)
European beech (*Fagus sylvatica*)
Evergreen oak (*Quercus ilex* L.)
Field maple (*Acer campestre*)
Ginkgo
Gleditsia L. spec.
Himalayan birch (*Betula utilis*)
Horse chestnut (*Aesculus hippocastanum*)
Italian alder (*Alnus cordata*)
Large-leaved linden (*Tilia platyphyllos*)
London plane (*Platanus hybrida*)
Magnolia sp.
Norway maple (*Acer platanoides*)
Pussy willow (*Salix caprea*)
Quercus L. spec.
Red gum (*Liquidambar styraciflua*)
Silver birch (*Betula pendula*)
Small-leaved linden (*Tilia cordata*)
Sycamore (*Acer pseudoplatanus*)
Tulip tree (*Liriodendron tulipifera* L.)
White oak (*Quercus pubescens*)

Alder buckthorn (*Rhamnus frangula*)
Blackthorn (*Prunus spinosa*)
Cherry laurel (*Prunus laurocerasus*)
Chokeberry (*Aronia prunifolia*)
Common box (*Buxus sempervirens*)
Common holly (*Ilex aquifolium* L.)
Common lilac (*Syringa vulgaris*)
Dogwoods (*Cornus spectabilis*)
Elaeagnus sp.
European hazel (*Corylus avellana*)
Forsythia (*Forsythia x intermedia*)
Garden privet (*Ligustrum ovalifolium*)
Golden currant (*Ribes aureum*)
Hibiscus L. spec.
Holly-leaved barberry (*Mahonia aquifolium*)
Japanese barberry (*Berberis thunbergii*)
Juneberry (*Amelanchier canadensis*)
Privets (*Ligustrum spectabilis*)
Prunus ornamental species
Prunus sp.
Red-flowered currant (*Ribes sanguineum*)
Rhododendron L. spec.
shrubby cinquefoil (*Potentilla fruticosa*)
Siberian pea tree (*Caragana arborescens*)
Snowberry (*Symphoricarpos*)
Spindle (*Euonymus europaeus*)
Spiraea sp.
Spiraea x vanhouttei
Viburnum tinus L.
White beech (*Carpinus betulus*)
Wild privet (*Ligustrum vulgare*)
Wintercreeper (*Euonymus fortunei*)

Austrian pine (*Pinus austriaca*)
Chinese juniper (*Juniperus media*)
Chinese thuja (*Thuja orientalis*)
Colorado spruce (*Picea pungens*)
Lawson's false cypress (*Chamaecyparis lawsoniana*)
Leyland cypress (*Cupressocyparis leylandii*)
Nordmann fir (*Abies nordmanniana*)
Northern white cedar (*Thuja occidentalis*)
Norway spruce (*Picea abies*)
Picea sp.
Scots pine (*Pinus sylvestris*)

Western red cedar Excelsa (*Thuja plicata excelsa*)

List of sensitive ornamental plants: spray drift may cause significant damage, such as necrosis, discoloration, chlorosis or stunting of European yew (*Taxus baccata*), *rosa*, *cotoneaster* and *crataegus* species.

Do not apply Valdor Flex around or under shrubs of the Rosaceae family.

Application around or under other species not listed here is not recommended.

SITUATION SPECIFIC INFORMATION

Use Areas

Natural surfaces not intended to bear vegetation, permeable surfaces overlying soil
Valdor Flex may be used in non-crop areas against weeds in open soil and against weeds growing in gravel or other porous surfaces. Examples of suitable use areas include gravel paths and driveways, porous surfaces alongside roadways and fence lines, porous strips of land adjacent to buildings, industrial sites, timber yards, farm yards, oil and gas storage sites, power stations, electric sub-stations, beneath pylons, around street/park obstacles and furniture, porous areas near to trees and shrubs and other natural surfaces where vegetation is not desirable.

Hard surfaces (railway ballast only)

Valdor Flex may be used on railway track, railway sidings and other ballast areas of rail infrastructure.

Amenity vegetation (around)

Valdor Flex may be used around the base of trees, shrubs and other plants in areas of semi-natural or ornamental vegetation, including parks, roadsides and other amenity areas. A drift shield is recommended for use around amenity vegetation, and in other areas where desired vegetation may be present.

General

Valdor Flex must not be used on non-porous man made surfaces, for example paved areas, concrete or tarmac car parks and footpaths.
Valdor Flex may be used on porous surfaces such as gravel ONLY where the underlying surface is soil. Do not use if an impermeable membrane lies between the gravel and the soil. The product must not be used on gravel where the underlying surface is concrete, tarmac or any other non-porous surface, or in situations where there is potential for run-off into surface waters.

Rates of Use

EQUIPMENT	AREA	PRODUCT REQUIRED	WATER VOLUME	Spray Quality (Nozzle)	Buffer Zone
Knapsack	200 m ²	1 sachet (10 g)	6 – 10 L	Coarse *Hypro Polijet ANO.6 or similar	2 m
Vehicle mounted sprayer	200 m ²	1 sachet (10 g)	6 – 10 L	Three-star drift reduction nozzles	6 m

* Hypro Polijet ANO.6 nozzle provided gives, subject to calibration: a coarse spray with a flow rate of 0.6 L/min at 1 bar, giving 225 L/ha at 4kph walking speed, swath width of 40 cm, 40 cm nozzle height.

Application Timing

Apply at any time of the year to weed-free soil (latest 2-leaf stage of any weeds present). Application to frozen ground should be avoided. 24 hours of dry weather are required immediately following application for optimum control.

MIXING AND SPRAYING

Half fill the spray tank with clean water and start gentle agitation. Add the required quantity of Valdor Flex. Top up to the required volume with water and agitate to ensure the granules are dissolved. Use immediately.

WASH OUT THE SPRAYER THOROUGHLY AFTER USE, USING A WETTING AGENT OR PROPRIETARY TANK CLEANER WITH TWO RINSES, AS TRACES OF VALDOR FLEX MAY CAUSE HARM TO OTHER SUSCEPTIBLE PLANTS SPRAYED LATER.

Equipment

Hand-held (natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, amenity vegetation and railway ballast)

Use a knapsack sprayer or tank and lance fitted with a coarse nozzle using a pressure of around 1-2 bars to provide a coarse spray. Use of anti-drift nozzles or the use of a protector shield to avoid any drift is recommended.

NOZZLE: Hypro Polijet ANO.6 nozzle gives, subject to calibration: a coarse spray with a flow rate of 0.6 L/min at 1 bar, giving 225 L/ha at 4kph walking speed, swath width of 40 cm, 40 cm nozzle height, or use similar nozzles that give coarse spray. Good and even coverage of foliage and soil is essential for optimum activity.

Vehicle-mounted applications (natural surfaces not intended to bear vegetation, permeable surfaces overlying soil, amenity vegetation and railway ballast)

Use a vehicle-mounted boom sprayer fitted with three-star drift reducing nozzles and a pressure of around 1-2 bars to provide a coarse spray. Good and even coverage of foliage and soil is essential for optimum activity.

COMPATIBILITY

Valdor Flex may be tank-mixed with other plant protection products providing that the application timing is correct for both Valdor Flex and the partner(s) in the mixture. For further information on the authorisation status of mixture partners, consult the manufacturer.

For 24 hour emergency information contact Bayer CropScience Limited
Telephone: 00800 1020 3333 or nearest National Poisons Information Centre

Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge
CB4 0WB
Telephone: 00800 1214 9451
www.environmentalscience.bayer.co.uk for SDS and larger label
© Registered trademark of Bayer CropScience Ltd.

GB86218453C - ARTICLE 85374958 - P2

Bayer